

# Boxwood-Phytophthora Root and Crown Rot

*Dr. Fulya Baysal-Gurel, Md Niamul Kabir and  
Angelo Randaci*

*Otis L. Floyd Nursery Research Center  
College of Agriculture  
Tennessee State University  
[fbaysalg@tnstate.edu](mailto:fbaysalg@tnstate.edu)*

**ANR-PATH-9-2017**

Phytophthora root and crown rot is caused by several species of Oomycetes (known as water molds). The *Phytophthora* species are *P. cinnamomi*, *P. nicotianae*, *P. citrophthora* and *P. occultans*. American (common), Japanese and English boxwood species are susceptible to this disease.

The pathogen can infect all growth stages of field or container grown boxwood plants. Warm, humid and rainy conditions favor disease development. *Phytophthora* spp. can survive in soil or potting mix as resting spores (oospores, chlamydospores), or inside infected plant tissues as mycelium for long periods, and can infect other healthy boxwoods or other host plants. When the environmental conditions are favorable, chlamydospores and oospores produce new mycelium or sporangia. Sporangia release zoospores into soil, and the zoospores swim toward healthy roots through saturated soil or irrigation water.

## **Symptoms**

The general symptoms include wilting, stunting, leaf shed, limb dieback, leaf chlorosis, off-color foliage, crown rot (gray to brown color) (Figure 1) and root rot (roots are brown and water-soaked). Depending on the disease progression in roots, foliar symptoms can be observed on a few branches or the entire plant (Figure 2).



Figure 1 and 2. Symptoms of boxwood *Phytophthora* crown and root rot.

### **Disease management**

*Phytophthora* root and crown rot is stimulated by rising soil temperature and moisture, poor drainage, excessive irrigation, extended periods of heavy rainfall or when plants are planted too deeply. Management of *Phytophthora* requires an integrated approach. Scouting and early diagnosis along with sound cultural practices such as sanitation, good drainage, proper irrigation, irrigation water treatments, crop rotation and chemical control are effective control strategies. Sanitize tools, equipment and reused pots (or avoid reusing pots altogether). If the production area or field had *Phytophthora* issues in the past, boxwood or other susceptible hosts should not be planted at least for three years.

Fungicides need to be applied before the establishment of the pathogen. A spray program that includes fungicides with different modes of action and FRAC codes is ideal for fungicide resistance management (Table 1).

If you would like to confirm that boxwood *Phytophthora* has infected your plants, you can submit a sample to your local university's plant diagnostic laboratory.

**Table 1.** List of selected fungicides and biorational products that can be used to prevent boxwood *Phytophthora* root and crown rot.

<b>Fungicide</b>	<b>Active ingredient</b>	<b>FRAC code</b>
Aliette WDG	Aluminum tris (O-ethyl phosphonate)	33
Alude	Mono and di-potassium salts of phosphorous acid	33
Adorn	Fluopicolide	43
Areca	Aluminum tris (O-ethyl phosphonate)	33
Banol	Propamocarb hydrochloride	28
Banrot	Etridiazole + thiophanate-methyl	1+14
Disarm O	Fluoxastrobin	11
Empress Intrinsic	Pyraclostrobin	11
Fosphite	Mono and di-potassium salts of phosphorous acid	33
Hurricane	Fludioxonil + mefenoxam	12 + 4
Mefenoxam 2 AQ	Mefenoxam	4
Micora	Mandipropamid	40
Orkestra	Fluxapyroxad + pyraclostrobin	7 + 11
Orvego	Ametoctradin + dimethomorph	45 + 40
OxiPhos	Mono and di-potassium salts of phosphorous acid + hydrogen peroxide	33
Pageant Intrinsic	Pyraclostrobin + boscalid	7 + 11
Root Shield	<i>Trichoderma harzianum</i> Rifai strain KRL-AG2	
RootShield PLUS+	<i>T. harzianum</i> strain T-22, and <i>T. virens</i> strain G-41	
Segovis	Oxathiapiprolin	U15
Segway O	Cyazofamid	21
Subdue GR	Mefenoxam	4
Subdue MAXX	Mefenoxam	4
Terrazole	Etridiazole	14
Truban	Etridiazole	14

**NOTE:** Before applying ANY disease management product, be sure to: 1) read the label to be sure that the product is allowed for the crop and the disease you intend to control; 2) read and understand the safety precautions and application restriction.

**For additional information, contact your local nursery specialist office at:**

**Tennessee State University**  
College of Agriculture  
3500 John A. Merritt Blvd., Box 9635 Nashville, TN 3720-1561  
<http://www.tnstate.edu/extension>

**Tennessee State University, Otis L. Floyd Nursery Research Center**  
472 Cadillac Lane McMinnville, TN 37110 <http://www.tnstate.edu/agriculture/nrc/>

#### **Precautionary Statement**

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label.

#### **Disclaimer**

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication. Use of trade, brand, or active ingredient names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar and suitable composition, nor does it guarantee or warrant the standard of the product. The author(s) and Tennessee State University assume no liability resulting from the use of these recommendations.

Dr. Chandra Reddy, Dean, Tennessee State University, College of Agriculture, Human and Natural Sciences  
Dr. Latif Lighari, Associate Dean Extension, Tennessee State University, College of Agriculture, Human and Natural Sciences  
Dr. Nick Gawel, Superintendent, Otis L. Floyd Nursery Research Center, Tennessee State University, College of Agriculture, Human and Natural Sciences

---

**TSU-18-0043(B)-15i-61605** Tennessee State University does not discriminate against students, employees, or applicants for admission or employment on the basis of race, color, religion, creed, national origin, sex, sexual orientation, gender identity/expression, disability, age, status as a protected veteran, genetic information, or any other legally protected class with respect to all employment, programs and activities sponsored by Tennessee State University. The following person has been designated to handle inquiries regarding non-discrimination policies: Rita Williams Seay, Office of Equity and Inclusion, [rseay@tnstate.edu](mailto:rseay@tnstate.edu), 3500 John Merritt Blvd., McWherter Administration Building, Suite 260, Nashville, TN 37209, 615-963-7438. The Tennessee State University policy on nondiscrimination can be found at [www.tnstate.edu/nondiscrimination](http://www.tnstate.edu/nondiscrimination).